

SCALED ASYMPTOTICS FOR SOME q -SERIES

RUIMING ZHANG

ABSTRACT. In this work we investigate the asymptotics for Euler's q -Exponential $E_q(z)$, q -Gamma function $\Gamma_q(z)$, Ramanujan's function $A_q(z)$, Jackson's q -Bessel function $J_\nu^{(2)}(z;q)$ of second kind, Stieltjes-Wigert orthogonal polynomials $S_n(x;q)$ and q -Laguerre polynomials $L_n^{(\alpha)}(x;q)$ as q approaching 1.

1. INTRODUCTION

Euler's q -Exponential $E_q(z)$, q -Gamma function $\Gamma_q(z)$, Ramanujan's function $A_q(z)$, Jackson's q -Bessel function $J_\nu^{(2)}(z;q)$ of second kind, Stieltjes-Wigert orthogonal polynomials $S_n(x;q)$ and q -Laguerre polynomials $L_n^{(\alpha)}(x;q)$ are very important examples in q -series, [5, 8, 11, 18]. They are used widely in other branches of mathematics and physics. In this work we will present a method to derive asymptotic formulas for these functions as q approaching 1 via Jacobi theta functions.

For any complex number a and parameter $0 < q < 1$, we define [5, 8, 11, 18]

$$(1) \quad (a; q)_\infty := \prod_{k=0}^{\infty} (1 - aq^k)$$

and the q -shifted factorial of a is defined by

$$(2) \quad (a; q)_n := \frac{(a; q)_\infty}{(aq^n; q)_\infty}, \quad n \in \mathbb{Z}.$$

We also use the following short-hand notation

$$(3) \quad (a_1, \dots, a_m; q)_n = \prod_{k=1}^m (a_k; q)_n, \quad m \in \mathbb{N}, \quad a_1, \dots, a_m \in \mathbb{C}, \quad n \in \mathbb{Z}.$$

Lemma 1.1. *Given any complex number a , assume that*

$$(4) \quad 0 < \frac{|a|q^n}{1-q} < \frac{1}{2}$$

for some positive integer n . Then,

$$(5) \quad \frac{(a; q)_\infty}{(a; q)_n} = (aq^n; q)_\infty := 1 + r_1(a; n)$$

with

$$(6) \quad |r_1(a; n)| \leq \frac{2|a|q^n}{1-q}$$

Date: November 30, 2006.

1991 *Mathematics Subject Classification.* Primary 30E15. Secondary 33D45.

and

$$(7) \quad \frac{(a; q)_n}{(a; q)_\infty} = \frac{1}{(aq^n; q)_\infty} := 1 + r_2(a; n)$$

with

$$(8) \quad |r_2(a; n)| \leq \frac{2|a|q^n}{(1-q)}.$$

Proof. From the q -binomial theorem [5, 8, 11, 18]

$$(9) \quad \frac{(az; q)_\infty}{(z; q)_\infty} = \sum_{k=0}^{\infty} \frac{(a; q)_k}{(q; q)_k} z^k \quad a, z \in \mathbb{C},$$

and the inequality

$$(10) \quad (q; q)_k \geq (1-q)^k$$

for $k = 0, 1, \dots$, we obtain

$$(11) \quad r_2(a; n) = \sum_{k=0}^{\infty} \frac{(aq^n)^{k+1}}{(q; q)_{k+1}}$$

and

$$(12) \quad |r_2(a; n)| \leq \sum_{k=0}^{\infty} \frac{(|a|q^n)^{k+1}}{(q; q)_{k+1}} \leq \frac{|a|q^n}{(1-q)} \sum_{k=0}^{\infty} \left(\frac{|a|q^n}{1-q} \right)^k \leq \frac{2|a|q^n}{(1-q)}.$$

Apply a limiting case of (9),

$$(13) \quad (z; q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q; q)_k} (-z)^k \quad z \in \mathbb{C},$$

and the inequalities,

$$(14) \quad \frac{1-q^k}{1-q} \geq kq^{k-1}, \quad \frac{(q; q)_k}{(1-q)^k} \geq k!q^{k(k-1)/2}, \quad \text{for } k = 0, 1, \dots$$

to obtain

$$(15) \quad r_1(a; n) = \sum_{k=1}^{\infty} \frac{q^{k(k-1)/2} (-aq^n)^k}{(q; q)_k},$$

and

$$(16) \quad \begin{aligned} |r_1(a; n)| &\leq \sum_{k=0}^{\infty} \frac{(|a|q^n)^{k+1}}{(1-q)^{k+1}} \frac{(1-q)^{k+1} q^{k(k+1)/2}}{(q; q)_{k+1}} \\ &\leq \sum_{k=0}^{\infty} \frac{(|a|q^n)^{k+1}}{(1-q)^{k+1}} \frac{1}{(k+1)!} \leq \frac{|a|q^n}{1-q} \exp(1/2) < \frac{2|a|q^n}{1-q}. \end{aligned}$$

□

The Dedekind $\eta(\tau)$ is defined as [22]

$$(17) \quad \eta(\tau) := e^{\pi i \tau / 12} \prod_{k=1}^{\infty} (1 - e^{2\pi i k \tau}),$$

or

$$(18) \quad \eta(\tau) = q^{1/12} (q^2; q^2)_\infty, \quad q = e^{\pi i \tau}, \quad \Im(\tau) > 0.$$

It has the transformation formula

$$(19) \quad \eta\left(-\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}} \eta(\tau).$$

The Jacobi theta functions are defined as

$$(20) \quad \theta_1(z; q) := \theta_1(v|\tau) := -i \sum_{k=-\infty}^{\infty} (-1)^k q^{(k+1/2)^2} e^{(2k+1)\pi i v},$$

$$(21) \quad \theta_2(z; q) := \theta_2(v|\tau) := \sum_{k=-\infty}^{\infty} q^{(k+1/2)^2} e^{(2k+1)\pi i v},$$

$$(22) \quad \theta_3(z; q) := \theta_3(v|\tau) := \sum_{k=-\infty}^{\infty} q^{k^2} e^{2k\pi i v},$$

$$(23) \quad \theta_4(z; q) := \theta_4(v|\tau) := \sum_{k=-\infty}^{\infty} (-1)^k q^{k^2} e^{2k\pi i v},$$

where

$$(24) \quad z = e^{2\pi i v}, \quad q = e^{\pi i \tau}, \quad \Im(\tau) > 0.$$

The Jacobi's triple product identities are

$$(25) \quad \theta_1(v|\tau) = 2q^{1/4} \sin \pi v (q^2; q^2)_{\infty} (q^2 e^{2\pi i v}; q^2)_{\infty} (q^2 e^{-2\pi i v}; q^2)_{\infty},$$

$$(26) \quad \theta_2(v|\tau) = 2q^{1/4} \cos \pi v (q^2; q^2)_{\infty} (-q^2 e^{2\pi i v}; q^2)_{\infty} (-q^2 e^{-2\pi i v}; q^2)_{\infty},$$

$$(27) \quad \theta_3(v|\tau) = (q^2; q^2)_{\infty} (-q e^{2\pi i v}; q^2)_{\infty} (-q e^{-2\pi i v}; q^2)_{\infty},$$

$$(28) \quad \theta_4(v|\tau) = (q^2; q^2)_{\infty} (q e^{2\pi i v}; q^2)_{\infty} (q e^{-2\pi i v}; q^2)_{\infty}.$$

The Jacobi θ functions satisfy transformations:

$$(29) \quad \theta_1\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = -i \sqrt{\frac{\tau}{i}} e^{\pi i v^2 / \tau} \theta_1(v \mid \tau),$$

$$(30) \quad \theta_2\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}} e^{\pi i v^2 / \tau} \theta_4(v \mid \tau),$$

$$(31) \quad \theta_3\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}} e^{\pi i v^2 / \tau} \theta_3(v \mid \tau),$$

$$(32) \quad \theta_4\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}} e^{\pi i v^2 / \tau} \theta_2(v \mid \tau).$$

Lemma 1.2. *For*

$$(33) \quad 0 < a < 1, \quad n \in \mathbb{N}, \quad \gamma > 0,$$

and

$$(34) \quad q = e^{-2\pi\gamma^{-1}n^{-a}},$$

we have

$$(35) \quad (q; q)_{\infty} = \sqrt{\gamma n^a} \exp\left\{\frac{\pi}{12}((\gamma n^a)^{-1} - \gamma n^a)\right\} \left\{1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\},$$

and

$$(36) \quad \frac{1}{(q; q)_{\infty}} = \frac{\exp\left\{\frac{\pi}{12}(\gamma n^a - (\gamma n^a)^{-1})\right\}}{\sqrt{\gamma n^a}} \left\{1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\}$$

as $n \rightarrow \infty$.

Proof. From formulas (17), (19) and (19) we get

$$\begin{aligned}
 (37) \quad (q; q)_\infty &= \exp(\pi\gamma^{-1}n^{-a}/12) \eta(\gamma^{-1}n^{-a}i) \\
 &= \sqrt{\gamma n^a} \exp(\pi\gamma^{-1}n^{-a}/12) \eta(\gamma n^a i) \\
 &= \sqrt{\gamma n^a} \exp(\pi\gamma^{-1}n^{-a}/12 - \pi\gamma n^a/12) \prod_{k=1}^{\infty} (1 - e^{-2\pi\gamma k n^a}) \\
 &= \sqrt{\gamma n^a} \exp(\pi\gamma^{-1}n^{-a}/12 - \pi\gamma n^a/12) \left\{ 1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right) \right\},
 \end{aligned}$$

and

$$(38) \quad \frac{1}{(q; q)_\infty} = \frac{\exp(\pi\gamma n^a/12 - \pi\gamma^{-1}n^{-a}/12)}{\sqrt{\gamma n^a}} \left\{ 1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right) \right\}$$

as $n \rightarrow \infty$. □

The Euler's q -Exponential is defined by [5, 8, 11, 18]

$$(39) \quad E_q(z) := (-z; q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q; q)_k} z^k, \quad z \in \mathbb{C}.$$

The q -Gamma function is defined as [5, 8, 11, 18]

$$(40) \quad \Gamma_q(x) := \frac{(q; q)_\infty}{(q^x; q)_\infty} (1 - q)^{1-x} \quad x \in \mathbb{C}.$$

Ramanujan's function $A_q(z)$ is defined as [11]

$$(41) \quad A_q(z) := \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q; q)_k} (-z)^k, \quad z \in \mathbb{C}.$$

Jackson's q -Bessel function of second kind [5, 8, 11, 18]

$$(42) \quad J_\nu^{(2)}(z; q) := \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{\nu+2k}}{(q, q^{\nu+1}; q)_k} q^{k(\nu+k)}, \quad \nu > -1, \quad z \in \mathbb{C}.$$

Stieltjes-Wigert orthogonal polynomials $\{S_n(x; q)\}_{n=0}^{\infty}$ are defined as [11]

$$(43) \quad S_n(x; q) := \sum_{k=0}^n \frac{q^{k^2} (-x)^k}{(q; q)_k (q; q)_{n-k}}, \quad x \in \mathbb{C}, \quad n \in \mathbb{N} \cup \{0\}.$$

Stieltjes-Wigert orthogonal polynomials come from an indeterminate moment problem. They satisfy the orthogonality relation

$$(44) \quad \int_0^\infty S_m(x; q) S_n(x; q) w_{sw}(x) dx = \frac{q^{-n}}{(q; q)_n} \delta_{m,n}, \quad n, m \in \mathbb{N} \cup \{0\},$$

where

$$(45) \quad w_{sw}(x) := \sqrt{\frac{-1}{2\pi \log q}} \exp\left(\frac{1}{2 \log q} \left[\log\left(\frac{x}{\sqrt{q}}\right)\right]^2\right), \quad x \in \mathbb{R}.$$

Clearly, the associated orthonormal Stieltjes-Wigert functions are given by

$$(46) \quad s_n(x; q) := \sqrt{q^n (q; q)_n w_{sw}(x)} S_n(x; q), \quad x \in \mathbb{R}^+, \quad n \in \mathbb{N} \cup \{0\}.$$

The q -Laguerre orthogonal polynomials $\{L_n^{(\alpha)}(x; q)\}_{n=0}^{\infty}$ are defined as [5, 8, 11, 18]

$$(47) \quad L_n^{(\alpha)}(x; q) := \sum_{k=0}^n \frac{q^{k^2+\alpha k} (-x)^k (q^{\alpha+1}; q)_n}{(q; q)_k (q, q^{\alpha+1}; q)_{n-k}}, \quad \alpha > -1, \quad x \in \mathbb{C}, \quad n \in \mathbb{N} \cup \{0\}.$$

The q -Laguerre orthogonal polynomials come from an indeterminate moment problem. They satisfy the orthogonality relation

$$(48) \quad \int_0^\infty L_m^{(\alpha)}(x; q) L_n^{(\alpha)}(x; q) w_{q\ell}(x) dx = \frac{(q^{\alpha+1}; q)_n}{q^n (q; q)_n} \delta_{m,n}, \quad \alpha > -1, \quad n, m \in \mathbb{N} \cup \{0\},$$

where

$$(49) \quad w_{q\ell}(x) := -\frac{\sin(\pi\alpha)}{\pi} \frac{(q; q)_\infty}{(q^{-\alpha}; q)_\infty} \frac{x^\alpha}{(-x; q)_\infty}, \quad x \in \mathbb{R}^+, \quad \alpha > -1.$$

Clearly, the associated orthonormal q -Laguerre orthogonal functions are given by

$$(50) \quad \ell_n(x; q) := \sqrt{\frac{q^n (q; q)_n}{(q^{\alpha+1}; q)_n}} w_{q\ell}(x) L_n^{(\alpha)}(x; q), \quad \alpha > -1, \quad n \in \mathbb{N} \cup \{0\}, \quad x \in \mathbb{R}^+.$$

For any positive integer n , we define

$$(51) \quad \chi(n) := 2 \left\{ \frac{n}{2} \right\},$$

then,

$$(52) \quad \chi(n) = n - 2 \left\lfloor \frac{n}{2} \right\rfloor = \left\lfloor \frac{n+1}{2} \right\rfloor - \left\lfloor \frac{n}{2} \right\rfloor,$$

where $\lfloor x \rfloor$ is the greatest integer less than or equals to $x \in \mathbb{R}$ and $\{x\}$ is the fractional part of $x \in \mathbb{R}$.

2. MAIN RESULTS

2.1. Euler q -Exponential Function $E_q(z)$, q -Gamma Function $\Gamma_q(x)$.

Theorem 2.1. *For*

$$(53) \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad u \in \mathbb{R}, \quad q = \exp(-2n^{-a}\pi),$$

we have

$$(54) \quad E_q(\exp 2\pi(u + n^{1-a} - \frac{1}{2}n^{-a})) = \exp \left\{ \pi n^{-a} (n^a u + n)^2 + \frac{\pi}{12} (n^a - n^{-a}) \right\} \left\{ 1 + \mathcal{O}(e^{-\pi n^a}) \right\},$$

and

$$(55) \quad E_q(-\exp 2\pi(u + n^{1-a} - \frac{1}{2}n^{-a})) = \frac{2 \exp(\pi n^{-a} (n^a u + n)^2) \cos(\pi n^a u)}{(-1)^n \exp \frac{\pi}{12} (2n^a - n^{-a})} \left\{ 1 + \mathcal{O}(e^{-2\pi n^a}) \right\}$$

as $n \rightarrow \infty$.

Similarly, $\Gamma_q(z)$ has asymptotic behavior:

Theorem 2.2. *For*

$$(56) \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad u \in \mathbb{R}, \quad q = \exp(-2n^{-a}\pi),$$

we have

$$(57) \quad \frac{1}{\Gamma_q\left(\frac{1}{2} - n - n^a u\right)} = \frac{2(-1)^n \exp(\pi n^{-a}(n^a u + n)^2) \cos(\pi n^a u) \{1 + \mathcal{O}(e^{-2\pi n^a})\}}{\sqrt{n^a} \exp(\pi n^a/12 + \pi n^{-a}/6) (1 - \exp(-2\pi n^{-a}))^{n+n^a u+1/2}},$$

and

$$(58) \quad \frac{1}{\Gamma_q\left(\frac{1}{2} + n + n^a u\right)} = \frac{\exp(\pi n^a/12 - \pi n^{-a}/12) \{1 + \mathcal{O}(e^{-2\pi n^a})\}}{\sqrt{n^a} (1 - e^{-2\pi n^{-a}})^{1/2 - n - n^a u}}$$

as $n \rightarrow \infty$.

2.2. Ramanujan's Function $A_q(z)$.

Theorem 2.3. *For*

$$(59) \quad 0 < q < 1, \quad z \in \mathbb{C} \setminus \{0\},$$

we have

$$(60) \quad A_q(q^{-2n}z) = \frac{(-z)^n \{\theta_4(z^{-1}; q) + e(n)\}}{(q; q)_\infty q^{n^2}},$$

and

$$(61) \quad |e(n)| \leq 4\theta_3(|z|^{-1}; q) \left\{ \frac{q^{n/2}}{1-q} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\}$$

for n sufficiently large.

Let

$$(62) \quad q = \exp(-\pi n^{-a}), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad u \in \mathbb{R},$$

then,

$$(63) \quad A_q(-\exp 2\pi(u + n^{1-a})) = \frac{\exp\{\pi n^{-a}(n^a u + n)^2\} \{1 + \mathcal{O}(\exp(-\pi n^a))\}}{\sqrt{2} \exp\{\pi n^{-a}/24 - \pi n^a/6\}},$$

and

$$(64) \quad A_q(\exp 2\pi(u + n^{1-a})) = \frac{\sqrt{2} \exp\{\pi n^{-a}(n^a u + n)^2\} \{\cos(\pi n^a u) + \mathcal{O}(\exp(-2\pi n^a))\}}{(-1)^n \exp \pi \{n^a/12 + n^{-a}/24\}}$$

as $n \rightarrow \infty$.

2.3. Jackson's q -Bessel function of second kind $J_\nu^{(2)}(z; q)$.

Theorem 2.4. *For*

$$(65) \quad 0 < q < 1, \quad z \in \mathbb{C} \setminus \{0\}, \quad \nu > -1,$$

we have

$$(66) \quad J_\nu^{(2)}\left(2\sqrt{zq^{-2n-\nu}}; q\right) = \frac{z^{n+\nu/2} \{\theta_4(z^{-1}; q) + e(n)\}}{(-1)^n (q; q)_\infty^2 q^{n^2+n\nu+\nu^2/2}},$$

and

$$(67) \quad |e(n)| \leq 12\theta_3(|z|^{-1}; q) \left\{ \frac{q^{n/2}}{1-q} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\}$$

for n sufficiently large.

Let

$$(68) \quad \nu > -1, \quad q = \exp(-\pi n^{-a}), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad u \in \mathbb{R},$$

then,

$$(69) \quad \begin{aligned} & J_\nu^{(2)} \left(2i \exp \left(\pi(u + n^{1-a} + \nu n^{-a}/2) \right); \exp(-\pi n^{-a}) \right) \\ &= \frac{i^\nu \exp \left\{ \pi n^{-a} (n^a u + n + \nu/2)^2 \right\} \{1 + \mathcal{O}(\exp(-\pi n^a))\}}{2\sqrt{n^a} \exp \pi \{n^{-a}/12 - n^a/3 - n^{-a}\nu^2/4\}}, \end{aligned}$$

and

$$(70) \quad \begin{aligned} & J_\nu^{(2)} \left(2 \exp \left(\pi(u + n^{1-a} + \nu n^{-a}/2) \right); \exp(-\pi n^{-a}) \right) \\ &= \frac{\exp \left\{ \pi n^{-a} (n^a u + n + \nu/2)^2 \right\} \{\cos(\pi n^a u) + \mathcal{O}(\exp(-2\pi n^a))\}}{(-1)^n \sqrt{n^a} \exp \pi \{n^{-a}/12 - n^a/12 - \nu^2 n^{-a}/4\}} \end{aligned}$$

as $n \rightarrow \infty$.

2.4. Stieltjes-Wigert Orthogonal Polynomials $S_n(x; q)$.

Theorem 2.5. *For*

$$(71) \quad 0 < q < 1, \quad z \in \mathbb{C} \setminus \{0\},$$

we have

$$(72) \quad S_n(zq^{-n}; q) = \frac{(-z)^{\lfloor n/2 \rfloor} \{\theta_4(z^{-1}q^{\chi(n)}; q) + e(n)\}}{(q; q)_\infty^2 q^{\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}},$$

and

$$(73) \quad |e(n)| \leq 12\theta_3(|z|^{-1} q^{\chi(n)}; q) \left\{ \frac{q^{n/4}}{1-q} + |z|^{\lfloor n/4 \rfloor} q^{\lfloor n/4 \rfloor^2 - \chi(n)\lfloor n/4 \rfloor} + \frac{q^{\lfloor n/4 \rfloor^2 + \chi(n)\lfloor n/4 \rfloor}}{|z|^{\lfloor n/4 \rfloor}} \right\}$$

for n sufficiently large.

Let

$$(74) \quad q = \exp(-2\pi n^{-a}), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad u \in \mathbb{R},$$

then,

$$(75) \quad S_n(-\exp 2\pi n^{-a}(n^a u + n); \exp(-2\pi n^{-a}))$$

$$(76) \quad = \frac{\exp \left\{ \pi n^{-a} (n^a u + n)^2/2 \right\} \{1 + \mathcal{O}(e^{-\pi n^a/2})\}}{\sqrt{2n^a} \exp \{ \pi n^{-a}/6 - \pi n^a/6 \}},$$

and

$$(77) \quad S_n \left(\exp \left(2\pi n^{-a} (n^a u + n) \right); \exp(-2\pi n^{-a}) \right) \\ = \sqrt{\frac{2}{n^a}} \frac{\exp \left\{ \frac{\pi n^{-a}}{2} (n^a u + n)^2 \right\} \left\{ \cos \frac{\pi}{2} (n^a u + n) + \mathcal{O}(e^{-\pi n^a}) \right\}}{\exp \{ \pi n^{-a} / 6 - \pi n^a / 24 \}}$$

for n sufficiently large. Consequently,

$$(78) \quad s_n(\exp 2\pi n^{-a} (n^a u + n); \exp(-2\pi n^{-a})) \\ = \frac{\exp(-u\pi/2) \left\{ \cos \frac{\pi}{2} (n^a u + n) + \mathcal{O}(e^{-\pi n^a}) \right\}}{\sqrt{\pi} \exp(3\pi n^{1-a}/2 + \pi n^{-a}/4)}$$

as $n \rightarrow \infty$.

2.5. q -Laguerre Orthogonal Polynomials $L_n^{(\alpha)}(x; q)$.

Theorem 2.6. For

$$(79) \quad 0 < q < 1, \quad z \in \mathbb{C} \setminus \{0\}, \quad \alpha > -1,$$

we have

$$(80) \quad L_n^{(\alpha)}(zq^{-n-\alpha}; q) = \frac{(-z)^{\lfloor n/2 \rfloor} \left\{ \theta_4(z^{-1}q^{\chi(n)}; q) + e(n) \right\}}{(q; q)_\infty^2 q^{\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}},$$

and

$$(81) \quad |e(n)| \leq 60\theta_3(|z|^{-1}q^{\chi(n)}; q) \left\{ \frac{q^{\lfloor n/4 \rfloor^2 + \chi(n)\lfloor n/4 \rfloor}}{|z|^{\lfloor n/4 \rfloor}} + |z|^{\lfloor n/4 \rfloor} q^{\lfloor n/4 \rfloor^2 - \chi(n)\lfloor n/4 \rfloor} + \frac{q^{n/4}}{1-q} \right\}$$

for sufficiently n . Let

$$(82) \quad q = \exp(-2\pi n^{-a}), \quad 0 < a < \frac{1}{2}, \quad \alpha > -1, \quad n \in \mathbb{N}, \quad u \in \mathbb{R},$$

then,

$$(83) \quad L_n^\alpha(-\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ = \frac{\exp \left\{ \frac{\pi n^{-a}}{2} (n^a u + n)^2 \right\} \left\{ 1 + \mathcal{O}(\exp(-\pi n^a/2)) \right\}}{\sqrt{2n^a} \exp \pi \{ \pi n^{-a} / 6 - \pi n^a / 6 \}},$$

and

$$(84) \quad L_n^\alpha(\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ = \sqrt{\frac{2}{n^a}} \frac{\exp \left\{ \frac{\pi n^{-a}}{2} (n^a u + n)^2 \right\} \left\{ \cos \frac{\pi}{2} (n^a u + n) + \mathcal{O}(\exp(-\pi n^a)) \right\}}{\exp \{ \pi n^a / 24 + \pi n^{-a} / 6 \}}$$

as $n \rightarrow \infty$. Consequently,

$$(85) \quad \ell_n^{(\alpha)}(\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ = \frac{1}{\sqrt{\pi}} \frac{\exp(-\pi u/2) \left\{ \cos \frac{\pi}{2} (n^a u + n) + \mathcal{O}(\exp(-\pi n^a)) \right\}}{\exp \frac{\pi}{2} (3n^{1-a} + \alpha^2 n^a + n^a/6 + 2\alpha n^{-a} + n^{-a}/2 - \alpha^2 n^{-a})}$$

as $n \rightarrow \infty$.

3. PROOFS

3.1. Proof for Theorem 2.1.

Proof. It is clear that

$$(86) \quad E_q(zq^{1/2-n}) = (-zq^{1/2-n}; q)_\infty = \frac{z^n q^{-n^2/2} (q, -zq^{1/2}, -z^{-1}q^{1/2}; q)_\infty}{(q, -z^{-1}q^{1/2+n}; q)_\infty}.$$

Thus,

$$(87) \quad \begin{aligned} E_q(q^{-n+1/2}e^{2\pi u}) &= (-q^{-n+1/2}e^{2\pi u}; q)_\infty \\ &= \frac{q^{-n^2/2}e^{2\pi nu} (q, -q^{1/2}e^{2\pi u}, -q^{1/2}e^{-2\pi u}; q)_\infty}{(q, -q^{n+1/2}e^{-2\pi u}; q)_\infty} \\ &= \frac{q^{-n^2/2}e^{2\pi nu}\theta_3(e^{2\pi u}; q^{1/2})}{(q, -q^{n+1/2}e^{-2\pi u}; q)_\infty}, \end{aligned}$$

and

$$(88) \quad \begin{aligned} E_q(-q^{-n+1/2}e^{2\pi u}) &= (q^{-n+1/2}e^{2\pi u}; q)_\infty \\ &= \frac{(-1)^n q^{-n^2/2}e^{2\pi nu} (q, q^{1/2}e^{2\pi u}, q^{1/2}e^{-2\pi u}; q)_\infty}{(q, q^{n+1/2}e^{-2\pi u}; q)_\infty} \\ &= \frac{(-1)^n q^{-n^2/2}e^{2\pi nu}\theta_4(e^{2\pi u}; q^{1/2})}{(q, q^{n+1/2}e^{-2\pi u}; q)_\infty}. \end{aligned}$$

Since

$$(89) \quad \frac{1}{(q; q)_\infty} = \frac{\exp\left\{\frac{\pi}{12}(n^a - n^{-a})\right\}}{\sqrt{n^a}} \left\{1 + \mathcal{O}\left(e^{-2\pi n^a}\right)\right\},$$

$$(90) \quad \frac{1}{(-q^{n+1/2}e^{-2\pi u}; q)_\infty} = 1 + \mathcal{O}\left(n^a e^{-2\pi n^{1-a}}\right),$$

$$(91) \quad \frac{1}{(q^{n+1/2}e^{-2\pi u}; q)_\infty} = 1 + \mathcal{O}\left(n^a e^{-2\pi n^{1-a}}\right),$$

$$(92) \quad \begin{aligned} \theta_4(e^{2\pi u}; q^{1/2}) &= \theta_4(ui|n^{-a}i) = \sqrt{n^a} \exp(\pi n^a u^2) \theta_2(n^a u|n^a i) \\ &= 2\sqrt{n^a} \exp\left(\pi n^a u^2 - \frac{\pi n^a}{4}\right) \cos(\pi n^a u) \left\{1 + \mathcal{O}\left(e^{-2\pi n^a}\right)\right\}, \end{aligned}$$

and

$$(93) \quad \begin{aligned} \theta_3(e^{2\pi u}; q^{1/2}) &= \theta_3(ui|n^{-a}i) = \sqrt{n^a} \exp(\pi n^a u^2) \theta_3(n^a u|n^a i) \\ &= \sqrt{n^a} \exp(\pi n^a u^2) \left\{1 + \mathcal{O}\left(e^{-\pi n^a}\right)\right\} \end{aligned}$$

as $n \rightarrow \infty$. Thus,

$$(94) \quad E_q(\exp 2\pi(u + n^{1-a} - \frac{1}{2}n^{-a})) = \exp\left\{\pi n^{-a}(n^a u + n)^2 + \frac{\pi}{12}(n^a - n^{-a})\right\} \left\{1 + \mathcal{O}\left(e^{-\pi n^a}\right)\right\},$$

and

(95)

$$E_q(-\exp 2\pi(u+n^{1-a}-\frac{1}{2}n^{-a})) = \frac{2 \exp(\pi n^{-a}(n^a u + n)^2) \cos(\pi n^a u)}{(-1)^n \exp \frac{\pi}{12}(2n^a - n^{-a})} \left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\}$$

as $n \rightarrow \infty$. $\square$

3.2. Proof for Theorem 2.2.

Proof. Observe that

$$(96) \quad \frac{1}{\Gamma_q(1/2 - n - n^a u)} = \frac{(q^{1/2-n} e^{2\pi u}; q)_\infty}{(q; q)_\infty (1-q)^{n+n^a u+1/2}} \\ = \frac{(q, q^{1/2} e^{-2\pi u}, q^{1/2} e^{2\pi u}; q)_\infty q^{-n^2/2} e^{2\pi n u}}{(-1)^n (1-q)^{n+n^a u+1/2} (q, q, q^{n+1/2} e^{-2\pi u}, q)_\infty}.$$

Since

$$(97) \quad \frac{1}{(q, q, q^{n+1/2} e^{-2\pi u}; q)_\infty} = n^{-a} \exp(\pi n^a/6 - \pi n^{-a}/6) \left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\},$$

and

$$(98) \quad (q, q^{1/2} e^{-2\pi u}, q^{1/2} e^{2\pi u}; q)_\infty = \theta_4(ui \mid n^{-a}i) = n^{a/2} e^{\pi n^a u^2} \theta_2(n^a u \mid n^a i) \\ = 2n^{a/2} \exp \pi n^a (u^2 - 1/4) \cos(n^a u \pi) \left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\}$$

as $n \rightarrow \infty$. Thus,

$$(99) \quad \frac{1}{\Gamma_q(1/2 - n - n^a u)} = \frac{2 \exp \pi (n^{-a}(n^a u + n)^2) \cos(\pi n^a u) \left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\}}{(-1)^n n^{a/2} (1 - e^{-2\pi n^{-a}})^{n+n^a u+1/2} \exp(\pi n^a/12 + \pi n^{-a}/6)}$$

as $n \rightarrow \infty$.

Similarly we could prove that

$$(100) \quad \frac{1}{\Gamma_q(1/2 + n + n^a u)} = \frac{\exp(\pi n^a/12 - \pi n^{-a}/12) \left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\}}{n^{a/2} (1 - e^{-2\pi n^{-a}})^{1/2-n-n^a u}}$$

as $n \rightarrow \infty$. $\square$

3.3. Proof for Theorem 2.3.

Proof. Write

$$(101) \quad A_q(q^{-2n} z)(q; q)_\infty = \sum_{k=0}^{\infty} \frac{(q; q)_\infty}{(q; q)_k} q^{k^2} (-z q^{-2n})^k \\ = \sum_{k=0}^n f(k) q^{k^2} (-z q^{-2n})^k + \sum_{k=n+1}^{\infty} f(k) q^{k^2} (-z q^{-2n})^k \\ = s_1 + s_2,$$

where

$$(102) \quad f(k) = (q^{k+1}; q)_\infty, \quad k \in \mathbb{N} \cup \{0\}.$$

Clearly,

$$(103) \quad |f(k)| \leq 1, \quad k \in \mathbb{N} \cup \{0\}.$$

Reverse summation order in s_1 to get

$$(104) \quad \frac{s_1 q^{n^2}}{(-z)^n} = \sum_{k=0}^n f(n-k) q^{k^2} (-z^{-1})^k.$$

By Lemma 1.1 we have

$$(105) \quad |f(n-k) - 1| = |r_1(q; n-k)| \leq \frac{2q^{n/2}}{1-q}$$

for $0 \leq k \leq \lfloor \frac{n}{2} \rfloor$ and n sufficiently large. Hence,

$$(106) \quad \begin{aligned} \frac{s_1 q^{n^2}}{(-z)^n} &= \sum_{k=0}^n f(n-k) q^{k^2} (-z^{-1})^k \\ &= \sum_{k=0}^{\infty} q^{k^2} (-z^{-1})^k - \sum_{k=\lfloor n/2 \rfloor}^{\infty} q^{k^2} (-z^{-1})^k \\ &\quad + \sum_{k=0}^{\lfloor n/2 \rfloor - 1} q^{k^2} (-z^{-1})^k \{f(n-k) - 1\} + \sum_{k=\lfloor n/2 \rfloor}^n q^{k^2} (-z^{-1})^k f(n-k) \\ &= \sum_{k=0}^{\infty} q^{k^2} (-q^{\chi(n)} z^{-1})^k + s_{11} + s_{12} + s_{13}. \end{aligned}$$

Thus,

$$(107) \quad |s_{11} + s_{13}| \leq 2 \sum_{k=\lfloor n/2 \rfloor}^{\infty} q^{k^2} \left(\frac{1}{|z|} \right)^k \leq 2 \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \sum_{k=0}^{\infty} q^{k^2} \left(\frac{1}{|z|} \right)^k \leq 2\theta_3(|z|^{-1}; q) \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}}$$

and

$$(108) \quad |s_{12}| \leq \frac{2q^{n/2}}{1-q} \sum_{k=0}^{\infty} q^{k^2} \left(\frac{1}{|z|} \right)^k \leq \frac{2\theta_3(|z|^{-1}; q)}{1-q} q^{n/2}$$

for n sufficiently large. Let

$$(109) \quad e_1(n) = s_{11} + s_{12} + s_{13},$$

then,

$$(110) \quad |e_1(n)| \leq 2\theta_3(|z|^{-1}; q) \left\{ \frac{q^{n/2}}{1-q} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\}$$

and

$$(111) \quad \frac{s_1 q^{n^2}}{(-z)^n} = \sum_{k=0}^{\infty} q^{k^2} (-z^{-1})^k + e_1(n),$$

for n sufficiently large.

Shift the summation index from k to $k+n$ in s_2 we obtain

$$(112) \quad \frac{s_2 q^{n^2}}{(-z)^n} = \sum_{k=1}^{\infty} q^{k^2} (-z)^k f(n+k).$$

Lemma 1.1 implies that

$$(113) \quad |f(n+k) - 1| = |r_1(q; n+k)| \leq \frac{2q^n}{1-q}$$

for $k \in \mathbb{N}$ and n sufficiently large. Thus,

$$(114) \quad \begin{aligned} \frac{s_2 q^{n^2}}{(-z)^n} &= \sum_{k=1}^{\infty} q^{k^2} (-z)^k f(n+k) \\ &= \sum_{k=1}^{\infty} q^{k^2} (-z)^k + \sum_{k=1}^{\infty} q^{k^2} (-z)^k \{f(n+k) - 1\} \\ &= \sum_{k=-1}^{-\infty} q^{k^2} (-z^{-1})^k + e_2(n), \end{aligned}$$

and

$$(115) \quad |e_2(n)| \leq \frac{2q^n}{1-q} \sum_{k=1}^{\infty} q^{k^2} |z|^k \leq \frac{2\theta_3(|z|^{-1}; q)}{1-q} q^{n/2}$$

for n sufficiently large. Hence,

$$(116) \quad \frac{A_q(q^{-2n}z)(q; q)_{\infty} q^{n^2}}{(-z)^n} = \theta_4(z^{-1}; q) + e(n),$$

with

$$(117) \quad |e(n)| \leq 4\theta_3(|z|^{-1}; q) \left\{ \frac{q^{n/2}}{1-q} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\}$$

for n sufficiently large.

From the properties of θ_3 and θ_4 we get

$$(118) \quad \begin{aligned} \theta_3(|z|^{-1}; q) &= \theta_3(iu \mid n^{-a}i) \\ &= \sqrt{n^a} \exp\{\pi n^a u^2\} \theta_3(n^a u \mid n^a i) \\ &= \sqrt{n^a} \exp\{\pi n^a u^2\} \left\{ 1 + \mathcal{O}(e^{-\pi n^a}) \right\}, \end{aligned}$$

and

$$(119) \quad \begin{aligned} \theta_4(z^{-1}; q) &= \theta_4(iu \mid n^{-a}i) \\ &= \sqrt{n^a} \exp\{\pi n^a u^2\} \theta_2(n^a u \mid n^a i) \\ &= 2\sqrt{n^a} \exp\{\pi n^a u^2 - \pi n^a/4\} \cos(\pi n^a u) \{1 + \mathcal{O}(\exp(-2\pi n^a))\} \end{aligned}$$

as $n \rightarrow \infty$. Observe that

$$(120) \quad n^a \ll n^{1-a}$$

for $0 < a < \frac{1}{2}$ as $n \rightarrow \infty$, thus,

$$(121) \quad A_q(-\exp 2\pi(u + n^{1-a})) = \frac{\exp\{\pi n^{-a}(n^a u + n)^2\} \{1 + \mathcal{O}(\exp(-\pi n^a))\}}{\sqrt{2} \exp\{\pi n^{-a}/24 - \pi n^a/6\}},$$

and

(122)

$$A_q(\exp 2\pi(u + n^{1-a})) = \frac{\sqrt{2} \exp \{ \pi n^{-a} (n^a u + n)^2 \} \{ \cos(\pi n^a u) + \mathcal{O}(\exp(-2\pi n^a)) \}}{(-1)^n \exp \pi \{ n^a/12 + n^{-a}/24 \}}$$

as $n \rightarrow \infty$. $\square$

3.4. Proof for Theorem 2.4.

Proof. It is clear that

$$(123) \quad \frac{J_\nu^{(2)}(2\sqrt{xq^{-\nu}}; q) (q; q)_\infty^2}{(xq^{-\nu})^{\nu/2}} = \sum_{k=0}^{\infty} q^{k^2} (-x)^k f(k),$$

where

$$(124) \quad f(k) := (q^{k+1}, q^{\nu+1+k}; q)_\infty, \quad k \in \mathbb{N} \cup \{0\}.$$

For $\nu > -1$, we have

$$(125) \quad |f(k)| \leq 1, \quad k \in \mathbb{N} \cup \{0\}.$$

It is clear that

$$(126) \quad \frac{J_\nu^{(2)}(2\sqrt{zq^{-2n-\nu}}; q) (q; q)_\infty^2 q^{n^2+n\nu+\nu^2/2}}{(-1)^n z^{n+\nu/2}} = \sum_{k=0}^n q^{k^2} (-z^{-1})^k f(n-k) + \sum_{k=1}^{\infty} q^{k^2} (-z)^k f(n+k).$$

Notice that for $k \geq 0$,

(127)

$$f(k) - 1 = \{r_1(q; k) + 1\} \{r_1(q^{\nu+1}; k) + 1\} - 1 = r_1(q; k) + r_1(q^{\nu+1}; k) + r_1(q; k)r_1(q^{\nu+1}; k),$$

then, for sufficiently large n ,

$$(128) \quad 0 < \frac{2q^{n/2}}{1-q} < 1,$$

and Lemma 1.1 implies

$$(129) \quad |f(n-k) - 1| \leq \frac{6q^{n/2}}{1-q}, \quad 0 \leq k \leq \left\lfloor \frac{n}{2} \right\rfloor,$$

and

$$(130) \quad |f(n+k) - 1| \leq \frac{6q^n}{1-q} < \frac{6q^{n/2}}{1-q}, \quad k \in \mathbb{N}.$$

Similar to the proof of Theorem 2.3, we have

$$(131) \quad J_\nu^{(2)}(2\sqrt{zq^{-2n-\nu}}; q) = \frac{z^{n+\nu/2} \{ \theta_4(z^{-1}; q) + e(n) \}}{(-1)^n (q; q)_\infty^2 q^{n^2+n\nu+\nu^2/2}},$$

and

$$(132) \quad |e(n)| \leq 12\theta_3(|z|^{-1}; q) \left\{ \frac{q^{n/2}}{1-q} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\}$$

for n sufficiently large. The rest of the proof is very similar to the corresponding part for Theorem 2.3. $\square$

3.5. Proof for Theorem 2.5.

Proof. As in the proof for Theorem 2.3 we have

(133)

$$\frac{(q; q)_\infty^2 S_n(zq^{-n}; q)}{(-z)^{\lfloor n/2 \rfloor} q^{-\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} q^{k^2} (-z^{-1} q^{\chi(n)})^k f\left(\left\lfloor \frac{n}{2} \right\rfloor - k\right) + \sum_{k=0}^{\lfloor \frac{n+1}{2} \rfloor} q^{k^2} (-z q^{-\chi(n)})^k f\left(\left\lfloor \frac{n}{2} \right\rfloor + k\right),$$

where

$$(134) \quad f(k) = (q^{k+1}; q)_\infty (q^{n-k+1}; q)_\infty, \quad k = 0, 1, \dots, n.$$

Clearly

$$(135) \quad |f(k)| \leq 1, \quad k = 0, 1, \dots, n.$$

Observe that

$$(136) \quad f(k) - 1 = r_1(q; k) + r_1(q; n-k) + r_1(q; k)r_1(q; n-k),$$

and if

$$(137) \quad 0 < \frac{2q^{n/4}}{1-q} < 1$$

for n sufficiently large, then Lemma 1.1 gives

$$(138) \quad |f\left(\left\lfloor \frac{n}{2} \right\rfloor - k\right) - 1| \leq \frac{6q^{n/4}}{1-q}, \quad 0 \leq k \leq \left\lfloor \frac{n}{4} \right\rfloor - 1,$$

and

$$(139) \quad |f\left(\left\lfloor \frac{n}{2} \right\rfloor + k\right) - 1| \leq \frac{6q^{n/4}}{1-q}, \quad k \in \mathbb{N}.$$

Thus,

$$(140) \quad \frac{(q; q)_\infty^2 S_n(zq^{-n}; q)}{(-z)^{\lfloor n/2 \rfloor} q^{-\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}} = \theta_4(z^{-1} q^{\chi(n)}; q) + e(n),$$

where

(141)

$$\begin{aligned} e(n) = & - \sum_{k=\lfloor n/4 \rfloor}^{\infty} q^{k^2} (-z^{-1} q^{\chi(n)})^k + \sum_{k=0}^{\lfloor n/4 \rfloor - 1} q^{k^2} (-z^{-1} q^{\chi(n)})^k \left(f\left(\left\lfloor \frac{n}{2} \right\rfloor - k\right) - 1 \right) \\ & + \sum_{k=\lfloor n/4 \rfloor}^{\lfloor n/2 \rfloor} q^{k^2} (-z^{-1} q^{\chi(n)})^k f\left(\left\lfloor \frac{n}{2} \right\rfloor - k\right) - \sum_{k=\lfloor n/4 \rfloor}^{\infty} q^{k^2} (-z q^{-\chi(n)})^k \\ & + \sum_{k=1}^{\lfloor n/4 \rfloor - 1} q^{k^2} (-z q^{-\chi(n)})^k \left(f\left(\left\lfloor \frac{n}{2} \right\rfloor + k\right) - 1 \right) + \sum_{k=\lfloor n/4 \rfloor}^{\lfloor (n+1)/2 \rfloor} q^{k^2} (-z q^{-\chi(n)})^k f\left(\left\lfloor \frac{n}{2} \right\rfloor + k\right). \end{aligned}$$

Therefore,

$$(142) \quad S_n(zq^{-n}; q) = \frac{(-z)^{\lfloor n/2 \rfloor} \{ \theta_4(z^{-1} q^{\chi(n)}; q) + e(n) \}}{(q; q)_\infty^2 q^{\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}},$$

and

(143)

$$|e(n)| \leq 12\theta_3 \left(|z|^{-1} q^{\chi(n)}; q \right) \left\{ \frac{q^{n/4}}{1-q} + |z|^{\lfloor n/4 \rfloor} q^{\lfloor n/4 \rfloor^2 - \chi(n)\lfloor n/4 \rfloor} + \frac{q^{\lfloor n/4 \rfloor^2 + \chi(n)\lfloor n/4 \rfloor}}{|z|^{\lfloor n/4 \rfloor}} \right\}$$

for n sufficiently large.

From the relations

$$(144) \quad \left\lfloor \frac{n}{2} \right\rfloor = \frac{n - \chi(n)}{2}, \quad \left\lceil \frac{n+1}{2} \right\rceil = \frac{n + \chi(n)}{2}, \quad n \in \mathbb{N},$$

Lemma 1.2 and transformation formulas of theta functions we get

$$(145) \quad \begin{aligned} & S_n(-\exp 2\pi n^{-a}(n^a u + n); \exp(-2\pi n^{-a})) \\ &= \frac{\exp \left\{ \pi n^{-a}(n^a u + n)^2/2 \right\} \left\{ 1 + \mathcal{O}(e^{-\pi n^a/2}) \right\}}{\sqrt{2} n^a \exp \left\{ \pi n^{-a}/6 - \pi n^a/6 \right\}}, \end{aligned}$$

and

$$(146) \quad \begin{aligned} & S_n(\exp 2\pi n^{-a}(n^a u + n); \exp(-2\pi n^{-a})) \\ &= \sqrt{\frac{2}{n^a}} \frac{\exp \left\{ \frac{\pi n^{-a}}{2}(n^a u + n) \right\} \left\{ \cos \frac{\pi}{2}(n^a u + n) + \mathcal{O}(e^{-\pi n^a}) \right\}}{\exp \left\{ \pi n^{-a}/6 - \pi n^a/24 \right\}} \end{aligned}$$

as $n \rightarrow \infty$.

Lemma 1.1 and Lemma 1.2 also give us that

$$(147) \quad (q; q)_n = \frac{(q; q)_\infty}{(q^{n+1}; q)_\infty} = \sqrt{n^a} \exp \left(\pi n^{-a}/12 - \pi n^a/12 \right) \left\{ 1 + \mathcal{O}(e^{-2\pi n^a}) \right\},$$

hence

$$(148) \quad \sqrt{w_{sw}(\exp 2\pi(u + n^{1-a}))} = \frac{\sqrt[4]{n^a}}{\sqrt{2\pi}} \exp \left(-\frac{n^a \pi}{2}(u + n^{1-a} + n^{-a}/2)^2 \right),$$

thus

$$(149) \quad \begin{aligned} & \sqrt{q^n(q; q)_n w_{sw}(\exp 2\pi(u + n^{1-a}))} \\ &= \sqrt{\frac{n^a}{2\pi}} \exp \left(-\frac{n^{-a} \pi}{2}(n^a u + n)^2 - \frac{u\pi}{2} \right) \\ & \times \exp \left(-\frac{\pi n^{-a}}{12} - \frac{\pi n^a}{24} - \frac{3\pi n^{1-a}}{2} \right) \left\{ 1 + \mathcal{O}(e^{-2\pi n^a}) \right\} \end{aligned}$$

as $n \rightarrow \infty$. Then we get

$$(150) \quad \begin{aligned} & s_n(\exp 2\pi n^{-a}(n^a u + n); \exp(-2\pi n^{-a})) \\ &= \frac{\exp(-u\pi/2) \left\{ \cos \frac{\pi}{2}(n^a u + n) + \mathcal{O}(e^{-\pi n^a}) \right\}}{\sqrt{\pi} \exp(3\pi n^{1-a}/2 + \pi n^{-a}/4)} \end{aligned}$$

as $n \rightarrow \infty$. □

3.6. Proof for Theorem 2.6.

Proof. Write

$$(151) \quad \frac{L_n^{(\alpha)}(zq^{-n-\alpha}; q)(q; q)_\infty^2}{(-z)^{\lfloor n/2 \rfloor} q^{-\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}} = \sum_{k=0}^{\lfloor n/2 \rfloor} q^{k^2} (-z^{-1} q^{\chi(n)})^k g\left(\left\lfloor \frac{n}{2} \right\rfloor - k\right) + \sum_{k=0}^{\lfloor (n+1)/2 \rfloor} q^{k^2} (-zq^{-\chi(n)})^k g\left(\left\lfloor \frac{n}{2} \right\rfloor + k\right),$$

where

$$(152) \quad g(k) = \frac{(q^{k+1}; q)_\infty (q^{n-k+1}; q)_\infty (q^{\alpha+n-k+1}; q)_\infty}{(q^{\alpha+n+1}; q)_\infty}, \quad k = 0, 1, \dots, n.$$

Then

$$(153) \quad |g(k)| \leq 1, \quad k = 0, 1, \dots, n.$$

Thus,

$$(154) \quad \frac{L_n^{(\alpha)}(zq^{-n-\alpha}; q)(q; q)_\infty^2}{(-z)^{\lfloor n/2 \rfloor} q^{-\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}} = \theta_4(z^{-1} q^{\chi(n)}; q) + e(n),$$

where

$$(155) \quad \begin{aligned} e(n) = & - \sum_{k=\lfloor n/4 \rfloor}^{\infty} q^{k^2} (-z^{-1} q^{\chi(n)})^k + \sum_{k=0}^{\lfloor n/4 \rfloor - 1} q^{k^2} (-z^{-1} q^{\chi(n)})^k (g(\lfloor \frac{n}{2} \rfloor - k) - 1) \\ & + \sum_{k=\lfloor n/4 \rfloor}^{\lfloor n/2 \rfloor} q^{k^2} (-z^{-1} q^{\chi(n)})^k g(\lfloor \frac{n}{2} \rfloor - k) - \sum_{k=\lfloor n/4 \rfloor}^{\infty} q^{k^2} (-zq^{-\chi(n)})^k \\ & + \sum_{k=0}^{\lfloor n/4 \rfloor - 1} q^{k^2} (-zq^{-\chi(n)})^k (g(\lfloor \frac{n}{2} \rfloor + k) - 1) + \sum_{k=\lfloor n/4 \rfloor}^{\lfloor (n+1)/2 \rfloor} q^{k^2} (-zq^{-\chi(n)})^k g(\lfloor \frac{n}{2} \rfloor + k). \end{aligned}$$

For $\alpha > -1$ and sufficiently large n with

$$(156) \quad 0 < \frac{2q^{n/4}}{1-q} < 1,$$

we expand

$$(157) \quad g(k) - 1 = \{r_2(q^{\alpha+1}; n) + 1\} \{r_1(q; n-k) + 1\} \{r_1(q; k) + 1\} \{r_1(q^{\alpha+1}; n-k) + 1\} - 1$$

and estimate each term using Lemma 1.1 to get

$$(158) \quad |g(\lfloor \frac{n}{2} \rfloor - k) - 1| \leq \frac{30q^{n/4}}{1-q}, \quad 0 \leq k \leq \lfloor \frac{n}{4} \rfloor,$$

and

$$(159) \quad |g(\lfloor \frac{n}{2} \rfloor + k) - 1| \leq \frac{30q^{n/2}}{1-q} < \frac{30q^{n/4}}{1-q}, \quad k \in \mathbb{N}.$$

Therefore,

$$(160) \quad L_n^{(\alpha)}(zq^{-n-\alpha}; q) = \frac{(-z)^{\lfloor n/2 \rfloor} \{\theta_4(z^{-1} q^{\chi(n)}; q) + e(n)\}}{(q; q)_\infty^2 q^{\lfloor n/2 \rfloor \lfloor (n+1)/2 \rfloor}},$$

and

(161)

$$|e(n)| \leq 60\theta_3(|z|^{-1}q^{\chi(n)}; q) \left\{ \frac{q^{\lfloor n/4 \rfloor^2 + \chi(n)\lfloor n/4 \rfloor}}{|z|^{\lfloor n/4 \rfloor}} + |z|^{\lfloor n/4 \rfloor} q^{\lfloor n/4 \rfloor^2 - \chi(n)\lfloor n/4 \rfloor} + \frac{q^{n/4}}{1-q} \right\}$$

for sufficiently n .

Formula (144) and transformations for theta function imply that

$$(162) \quad \begin{aligned} & L_n^\alpha(-\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ &= \frac{\exp\left\{\frac{\pi n^{-a}}{2}(n^a u + n)^2\right\} \{1 + \mathcal{O}(\exp(-\pi n^a/2))\}}{\sqrt{2n^a} \exp \pi \{\pi n^{-a}/6 - \pi n^a/6\}}, \end{aligned}$$

and

$$(163) \quad \begin{aligned} & L_n^\alpha(\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ &= \sqrt{\frac{2}{n^a}} \frac{\exp\left\{\frac{\pi n^{-a}}{2}(n^a u + n)^2\right\} \left\{\cos \frac{\pi}{2}(n^a u + n) + \mathcal{O}(\exp(-\pi n^a))\right\}}{\exp\{\pi n^a/24 + \pi n^{-a}/6\}} \end{aligned}$$

as $n \rightarrow \infty$.

Since

(164)

$$\begin{aligned} \frac{q^n(q; q)_n}{(q^{\alpha+1}; q)_n} w_{q^\ell}(zq^{-n-\alpha}) &= -\frac{\sin(\pi\alpha)(q; q)_\infty}{\pi(q^{-\alpha}; q)_\infty} \frac{z^\alpha q^{(1-\alpha)n-\alpha^2}(q; q)_n}{(q^{\alpha+1}; q)_n(-zq^{-n-\alpha}; q)_\infty} \\ &= -\frac{\sin(\pi\alpha)(q; q)_\infty^4}{\pi(1-q^{-\alpha})(q, q^{1-\alpha}, q^{\alpha+1}; q)_\infty} \\ &\quad \times \frac{z^{-n+\alpha} q^{n(n+3)/2-\alpha^2}}{(1+zq^{-\alpha})(q, -zq^{1-\alpha} - z^{-1}q^{1+\alpha}; q)_\infty} \\ &\quad \times \{1 + \mathcal{O}(n^a \exp(-2\pi n^{1-a}))\} \\ &= \frac{\exp \pi(u(2\alpha - 2n - 1) + 2\alpha^2 n^{-a} - n^{1-a}(n+3) - 2\alpha n^{-a} - n^{-a}/6 - n^a/3)}{\theta_1(\alpha n^{-a}i|n^{-a}i)\theta_2(ui + \alpha n^{-a}i|n^{-a}i)} \\ &\quad \times \frac{in^{2a} \sin \pi\alpha}{\pi} \{1 + \mathcal{O}(\exp(-2\pi n^a))\} \\ &= \frac{n^a \exp(-\pi n^{-a}(n^a u + n)^2 - \pi u) \{1 + \mathcal{O}(\exp(-\pi n^a))\}}{2\pi \exp \pi(3n^{1-a} + 2\alpha n^{-a} - \alpha^2 n^{-a} + \alpha^2 n^a + n^{-a}/6 + n^a/12)} \end{aligned}$$

for n sufficiently large. Therefore,

$$(165) \quad \begin{aligned} & \ell_n^{(\alpha)}(\exp 2\pi(u + n^{1-a} + \alpha n^{-a}); \exp(-2\pi n^{-a})) \\ &= \frac{1}{\sqrt{\pi}} \frac{\exp(-\pi u/2) \left\{\cos \frac{\pi}{2}(n^a u + n) + \mathcal{O}(\exp(-\pi n^a))\right\}}{\exp \frac{\pi}{2}(3n^{1-a} + \alpha^2 n^a + n^a/6 + 2\alpha n^{-a} + n^{-a}/2 - \alpha^2 n^{-a})} \end{aligned}$$

as $n \rightarrow \infty$. □

REFERENCES

- [1] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, English translation, Oliver and Boyd, Edinburgh, 1965.

- [2] G. E. Andrews, *q-series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra*, CBMS Regional Conference Series, number 66, American Mathematical Society, Providence, R.I. 1986.
- [3] G. E. Andrews, Ramanujan's "Lost" Note book VIII: The entire Rogers-Ramanujan function, *Advances in Math.* **191** (2005), 393–407.
- [4] G. E. Andrews, Ramanujan's "Lost" Note book IX: The entire Rogers-Ramanujan function, *Advances in Math.* **191** (2005), 408–422.
- [5] G. E. Andrews, R. A. Askey, and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.
- [6] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach*, American Mathematical Society, Providence, 2000.
- [7] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.* **52** (1999), 1491–1552.
- [8] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition Cambridge University Press, Cambridge, 2004.
- [9] W. K. Hayman, On the zeros of a q -Bessel function, *Contemporary Mathematics*, volume 382, American Mathematical Society, Providence, 2005, 205–216.
- [10] M. E. H. Ismail, Asymptotics of q -orthogonal polynomials and a q -Airy function, *Internat. Math. Res. Notices* 2005 No 18 (2005), 1063–1088.
- [11] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one Variable*, Cambridge University Press, Cambridge, 2005.
- [12] M. E. H. Ismail and X. Li, *Bounds for extreme zeros of orthogonal polynomials*, *Proc. Amer. Math. Soc.* **115** (1992), 131–140.
- [13] M. E. H. Ismail and D. R. Masson, q -Hermite polynomials, biorthogonal rational functions, *Trans. Amer. Math. Soc.* **346** (1994), 63–116.
- [14] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, *Advances in Math.*, (2007), to appear.
- [15] M. E. H. Ismail and R. Zhang, *Scaled Asymptotics for q -Polynomials*, *Comptes Rendus*, Vol. 344, Issue 2, 15 January 2007, Pages 71–75.
- [16] M. E. H. Ismail and R. Zhang, *Chaotic and Periodic Asymptotics for q -Orthogonal Polynomials*, joint with Mourad E.H. Ismail, *International Mathematics Research Notices*, Accepted.
- [17] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, Y. Yamada, Hypergeometric solutions to the q -Painlevé equations, *Internat. Math. Res. Notices* 47 (2004), 2497–2521.
- [18] R. Koekoek and R. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogues*, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.
- [19] M. L. Mehta, *Random Matrices*, third edition, Elsevier, Amsterdam, 2004.
- [20] W.-Y. Qiu and R. Wong, Uniform asymptotic formula for orthogonal polynomials with exponential weight, *SIAM J. Math. Anal.* **31** (2000), 992–1029.
- [21] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers* (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.
- [22] Hans Rademacher, *Topics in Analytic Number Theory*, Die Grundlehren der mathematischen Wissenschaften, Bd. 169, Springer-Verlag, New York-Heidelberg-Berlin, 1973. Z. 253. 10002.
- [23] G. Szegő, *Orthogonal Polynomials*, Fourth Edition, Amer. Math. Soc., Providence, 1975.
- [24] Z. Wang and R. Wong, Uniform asymptotics for the Stieltjes-Wigert polynomials: the Riemann-Hilbert approach, to appear.
- [25] R. Wong, *Asymptotic Approximations of Integrals*, Academic Press, Boston, 1989.
- [26] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, fourth edition, Cambridge University Press, Cambridge, 1927.

E-mail address: ruimingzhang@yahoo.com

Current address: School of Mathematics, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.

ADJUNCT PROFESSOR, BINZHOU VOCATIONAL COLLEGE, 533 BOHAI 9 ROAD, BINZHOU CITY, SHANDONG 256624, P. R. CHINA.